



INVESTOR IN PEOPLE


The Patent Office  
 Concept House  
 Cardiff Road  
 Newport  
 South Wales  
 NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.




*Stephen Hardley*

Signed

Dated 11 January 2005

BEST AVAILABLE COPY





The  
Patent  
Office

1/77

# Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road  
Newport  
South Wales  
NP9 1RH

1. Your reference P36233GB/KVC

11 DEC 2003

12DEC03 E858868-3 000056

FC1/7700 0-00-0328747.1 CHEQUE

2. Patent application number

(The Patent Office will fill in this part)

0328747.1

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Mars UK Limited  
3D Dundee Road, Slough  
Berkshire SL1 4LG

Patents ADP number (if you know it)

07475973001

If the applicant is a corporate body, give the country/state of its incorporation

United Kingdom

4. Title of the invention

Methods

5. Name of your agent (if you have one)

Kilburn & Strode  
20 Red Lion Street  
London  
WC1R 4PJ

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Patents ADP number (if you know it)

125001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number  
(if you know it)

Date of filing  
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing  
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
  - b) there is an inventor who is not named as an applicant, or
  - c) any named applicant is a corporate body.
- See note (d))

YES



## Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 10

Claim(s) 1

Abstract

Drawing(s) 3 + 5 *ll*

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 1/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents  
(please specify)

11. I/We request the grant of a patent on the basis of this application.

Signature

Date 11/12/03

*Kristina Cornish*

12. Name and daytime telephone number of person to contact in the United Kingdom

Kristina Cornish  
Tel: 020 7539 4200

### Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

### Notes

- a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.  
b) Write your answers in capital letters using black ink or you may type them.  
c) If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.  
d) If you have answered 'Yes' Patents Form 7/77 will need to be filed.  
e) Once you have filled in the form you must remember to sign and date it.  
f) For details of the fee and ways to pay please contact the Patent Office.



## Methods

Feline odontoclastic resorptive lesions (FORL) are one of the most common oral conditions in domestic cats, with prevalence increasing with age. During disease, odontoclasts mediate root resorption that is combined with the deposition of bone or cementum-like tissue by blast cells in an attempt to repair the damage. Root tissue is progressively lost, undermining the crown that may eventually break.

The prevalence of FORL in the domestic cat population has been reported to be 2-75%. The variation can be accounted for by differences in age, source of the population studied and diagnostic methods used.

Diagnosis of FORL is made by clinical examination (visual inspection and tactile exploration) or by a combination of clinical and radiographical examination. Clinical examination alone can only detect FORL above the gingival margin, missing lesions occurring in the root of the tooth and therefore radiographic studies are recommended. Radiographic changes are usually discrete and therefore clarity and detail are essential. For a dental radiograph to be diagnostic, it should be an accurate representation of the size and shape of the tooth without superimposition of adjacent structures. Therefore intraoral radiographic techniques are required; a parallel technique (Fig. 1) for the mandibular premolars and molars, and a bisecting angle technique (Fig. 2) for all other teeth. Whilst the parallel technique requires minimal radiographic training, the bisecting angle method is technically difficult and requires extensive training and experience, placing it beyond the reach of many veterinary practices.



'Full mouth radiographs' is the term used to describe a series of dental films where each tooth of the dentition is accurately depicted in at least one view. This is the gold-standard for FORL diagnosis, but requires 8-10 separate films that may take 15-30 minutes to obtain, during which time the subject must be  
 5 fully anaesthetised. The eight basic views (each view indicated in square bracket, below) are as follows:

Incisor view in the upper jaw (maxillary incisor teeth) [1]

Lateral view for each of the canines of the upper jaw (right maxillary canine and left maxillary canine) [2]

- 10 Left and right maxillary premolar and molar views (right maxillary premolar and molar, and left maxillary premolar and molar) [2]

Mandibular incisor and canine view (mandibular incisors and canines) [1]

Left and right mandibular premolar and molar views. (right mandibular premolar and molar, and left mandibular premolar and molar) [2]

15

In addition, lateral mandibular canine views (2 films) may be required.

It is disputed which teeth are most commonly affected with FORL. Ingham and others (2001), in Prevalence of odontoclastic resorptive lesions in a population of clinically healthy cats, *Journal of Small Animal Practice* 42, 439-443 have  
 20 reported in their study that the teeth most commonly affected with FORL were the third mandibular premolars. However, in a later study, Van Wessum, R., Harvey, C. E. & Hennet, P. (1992), Feline dental resorptive lesions, Prevalence patterns, *Veterinary Clinics of North America: Small Animal Practice* 22, 1405-1416 report that the mandibular and maxillary fourth  
 25 premolars are the most commonly affected with FORL. It was therefore of interest to determine if the FORL status of two teeth alone provided a reliable indicator of overall FORL status. The present study aimed to determine if FORL could be accurately diagnosed from the left and right mandibular



premolar and molar views (2 intraoral films and parallel radiographic technique) alone.

5 Accordingly, the present invention provides a method of diagnosing FORL, comprising taking no more than seven radiographic views of teeth from a feline animal, wherein at least one view is of each of the mandibular premolar teeth.

10 The invention comprises taking two, three, four, five, six or seven radiographic views of teeth from the mouth of a feline animal, wherein at least one view is of each of the mandibular third premolar teeth.

Preferably, at least two radiographic views are parallel views. Most preferably, these two parallel views are of the mandibular third premolar teeth.

15 The parallel technique is used usually for the mandibular premolars and molars. Details with regard to use the parallel technique is described herein with reference to Figure 1. For the other teeth, a bisecting angle technique is used. A discussion with regard to use of the bisecting angle technique is disclosed herein with reference to Figure 2.

20 It should be noted that oral examination, charting and radiology in accordance with the invention are usually (and may exclusively be) performed by a veterinary nurse and/or a dental technician whom are not veterinary specialists in dentistry. Accordingly, the radiographic results do not make it immediately  
25 possible to decide on a particular course of medical treatment.

The present invention particularly relates to a method of obtaining radiographic data of feline teeth, the method comprising taking no more than seven radiographic views of teeth from a mouth of a feline animal, wherein at least



one view is of each of the mandibular third premolar teeth. This method may comprise taking two, three, four, five, six or seven radiographic views of teeth from the mouth of a feline animal.

- 5 Preferably, at least two radiographic views are parallel views. Most preferably, these two parallel views are of the mandibular third premolar teeth.

- 10 The present invention also relates to the use of no more than seven radiographic views of teeth from the mouth of a feline animal, wherein at least one view is of each of the mandibular third premolar teeth, and the manufacturer for a tool for diagnosing FORL.

This use may comprise two, three, four, five or six radiographic views.

- 15 Preferably, at least two radiographic views are parallel. Most preferably, the two parallel views are of the mandibular premolar teeth.

- 20 The present invention also relates to dental treatment performed as the result of the diagnosis of FORL in the teeth of the feline animal, following a method according to the present invention. Accordingly, the method invention relates to a method as described herein according to the invention followed by the appropriate medical treatment.

- 25 The present invention relates to the number of radiographic views needed in order to determine an initial result for the presence or absence of FORL. If the initial less than seven radiographic views provide the result that FORL is present, it may be necessary or appropriate to take further radiographic views of additional teeth. Thus, after an initial result is obtained, further radiographic views, to a total of eight or over may be taken. However, the advantage of the



present invention is that less than eight views are initially required in order to determine whether further time, money and effort are required by taking additional views.

- 5 The present invention is described with reference to the figures, in which:

Figure 1 shows the Parallel Technique

- 10 The parallel technique is used for the mandibular premolars and the molars. The patient is placed in lateral recumbency (with the side to be radiographed uppermost). The film is placed between the tongue and the teeth and pushed as far down into the sublingual fossa as possible. The X-ray beam is then directed from lateral to medial at right angles to the long axis of the tooth. The resulting image of the tooth has very little magnification or distortion. Due to the anatomy of the oral cavity, this technique is only possible in the mandibular premolar and molar regions.

Figure 2 shows the bisecting angle technique

- 20 The Bisecting angle technique is required to minimise distortion when taking radiographs of the teeth in the upper jaw and at the mandibular incisors and canines. The film is positioned at an angle behind the tooth in question. To avoid foreshortening or elongation of the image, an imaginary plane is drawn half way between the plane of the film and a plane through the long axis of the tooth, i.e. at the bisecting angle, and the X-ray beam is directed perpendicular to this plane. In this way, both sides of the triangles formed are the same length and the resulting image of the tooth is similar to the real tooth.
- 25

Figure 3 shows the side view of a cat skull with teeth shown.

Figure 4 shows the numbering of the maxillary and mandibular teeth



The invention is now described with and referenced to the following example:

### *Example*

5

#### *Introduction*

10 The FORL status (presence or absence of odontoclastic resorptive lesions) of 423 clinically healthy cats was determined based on radiographic findings in a series of full mouth radiographs (8 views). This status was compared to FORL status based on evaluation of only two views, namely the right and left mandibular premolar and molar views. Using the FORL status of the right and left third mandibular premolars (307 and 407) alone correctly predicted overall FORL status in 93.4% cats. Overall FORL status can therefore be confidently  
15 diagnosed in 9/10 cats by assessing FORL status in just two teeth (307, 407) using two films, which has benefits for the cat (less anaesthetic time and reduced exposure to radiation) and owner (reduced cost of screening).

#### *Materials and Methods*

20

##### *Subjects*

Four hundred and twenty-three cats (mean age  $4.44 \pm \text{SEM } 0.14$  years, age range 1-14 years; 414 domestic shorthaired and 9 British Blue) were studied,  
25 consisting of 243 females (45 entire, 198 neutered) and 180 males (9 entire, 171 neutered). The cats were all bred and maintained by the WALTHAM Centre for Pet Nutrition, housed in groups of 20-25 in purpose-built, environmentally enriched surroundings. All cats had been fed a variety of dry and wet commercially available pet foods throughout their lives, and were

---



clinically healthy as determined by a full clinical physical examination. This study complies with Waltham's rigorous ethical policy.

### *Study Design*

5

FORL status was assessed on a single occasion. Each cat underwent a thorough oral examination and a full mouth series of dental radiographs were taken under general anaesthesia at the time of a scheduled dental cleaning. The oral examination, charting and radiography were performed by a listed  
10 veterinary nurse and a dental technician who had received training from, and worked under the supervision of, a European veterinary specialist in dentistry. Radiographic interpretation was performed by the specialist. Where indicated, additional dental treatment was also performed.

15

### *General Anaesthesia*

20

The cats were pre-medicated with a combination of 0.4 mg/kg butorphanol (Torbugesic; Fort Dodge Animal Health) and 0.05 mg/kg medetomidine (Domitor; Pfizer) by subcutaneous injection, and anaesthesia was induced by  
masking with 2 per cent isoflurane (IsoFlo Vet; Schering-Plough Animal Health) in oxygen. A cuffed endotracheal tube was placed and anaesthesia was maintained using isoflurane in oxygen and a non-breathing Ayre's T-piece anaesthetic circuit. The medetomidine sedation was reversed once treatment had been completed by the intramuscular injection of 0.125 mg/kg atipamezole  
25 (Antisedan; Pfizer).



*FORL status*

Each cat underwent a full clinical oral examination, including gingivitis scoring, and a full mouth series of dental radiographs were obtained. The oral  
5 examination consisted of visual inspection and tactile exploration of the tooth surfaces using a dental explorer. In addition, the periodontium was investigated using a blunt-ended periodontal probe.

Radiography consisted of eight views, using intra-oral placement of periapical  
10 size dental film and a dental x-ray machine. Mandibular premolar and molar views were obtained using a parallel technique, whilst all other views were taken using a bisecting angle technique.

The presence of FORL was determined by radiographic features such as loss of  
15 integrity of periodontal ligament space, irregularities in the root surface, radiolucent areas within the root dentine often extending into the crown dentine, and replacement of root substance by bone-like tissue. Teeth were classified as missing if they were absent on the radiograph. Some teeth were missing clinically, but the resorbing roots were present on the radiograph; these  
20 teeth were classified as teeth affected by FORL rather than missing. Teeth were also identified (306, 406) that are not recognised as part of the normal feline dentition and were disregarded from the data set.

*Model and Statistics*

25 Using the eight-film, full screening procedure, the true FORL status of each cat was determined. In addition, the test FORL status was determined using the results from 307, 407 on the two premolar and molar views alone.



Missing teeth could not be assumed to be missing due to FORL, and the following rationale was applied. If 307 and 407 were both present they were scored as usual. If 307 and 407 were both missing completely (and the resorbing roots were not radiographically visible), it could not be assumed that the loss was due to FORL and so this situation was scored as FORL absent. Where either 307 or 407 were missing, but there was no evidence of FORL in the other one, again this was scored FORL absent.

The true FORL status (8 film) was then compared to the test FORL status (2 film) from simple cross-tabulation tables.

### Results

The prevalence rate of FORL in the sample of cats was 30.7% (table 1). The true FORL status was correctly predicted using two films alone in 93.4% cats (table 1).

Table 1: Comparison of FORL status in 423 cats using two methods. True FORL status was determined using a full mouth (8 film) series, whilst test FORL status was determined using a reduced (2 film) series of dental radiographs.

Table 1

Number of cats (n=423)		True FORL status (8 film)	
		FORL +	FORL -
Test FORL status (2 film)	FORL +	102	0
	FORL -	28	293



### *Discussion*

The current study has shown that FORL status can be diagnosed in 9/10 cats using just two radiographs. This finding has important implications for both the cat, which is exposed to shorter anaesthesia and less radiation, and the veterinary practice, since the investigation can be conducted rapidly and at a reduced cost to the client. The technique is therefore suitable as a rapid screening procedure for FORL that can be adopted by all practices. These two views require intraoral positioning of dental film, but can be taken with a conventional veterinary radiographic unit.

15

The adoption of a simplified, standardised technique for FORL diagnosis would assist in the estimation of prevalence of the disease, which has been reported to be anywhere between 2-75%.

### *Conclusion*

20

This study provides a rapid screening technique for FORL that will confidently diagnose 9/10 cats and that can be adopted by all veterinary practices. The radiographic technique is easy to learn and does not require purchase of specialised equipment as these views can be taken with a routine veterinary X-ray unit. It is recommended that left and right mandibular premolar and molar views are taken of all cats anaesthetised for dental treatment.



## CLAIMS:

1. A method of diagnosing FORL comprising taking no more than seven radiographic views of teeth from the mouth of a feline animal, wherein at least  
5 one view is of each of the mandibular third premolar teeth.
2. A method, as claimed in claim 1, comprising taking no more than four radiographic views of teeth from the mouth of a feline animal.
- 10 3. A method, as claimed in claim 1 or 2, comprising taking no more than two radiographic views of teeth from the mouth of a feline animal.
4. A method, as claimed in any one of claims 1 to 3 wherein at least two radiographic views are parallel views.
- 15 5. Use of no more than seven radiographic views of teeth from the mouth of a feline animal, wherein at least one view is of each of the mandibular third premolar teeth, is the manufacture of a tool for diagnosing FORL.
- 20 6. A method of diagnosing FORL as herein before described with reference to the example.






Figure 1

5




Figure 2






Figure 3






Figure 4



# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/GB04/005201

International filing date: 10 December 2004 (10.12.2004)

Document type: Certified copy of priority document

Document details: Country/Office: GB  
Number: 0328747.1  
Filing date: 11 December 2003 (11.12.2003)

Date of receipt at the International Bureau: 24 January 2005 (24.01.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**